IN THE CLAIMS

This listing of claims will replace all prior versions, and listings, of claims in the

application.

Claims 1-16 (Canceled).

Claim 17 (Currently Amended): A method of stabilizing an emulsion comprising

adding, during preparation of the emulsion, mixture comprising a surfactant and a

cosurfactant, the cosurfactant being an amphiphilic polymer having one or more hydrophobic

subunits (A) and one or more hydrophilic subunits (B), wherein one or more of said

hydrophobic subunits (A) have been formed on the basis of a polyisobutene block whose

polyisobutene macromolecules have terminal double bonds to an extent of at least 50 mol%

in an amount sufficient to stabilize the emulsion.

Claim 18 (Currently Amended): The mixture method according to claim 17, wherein

every hydrophobic subunit (A) has been formed on the basis of a polyisobutene block whose

polyisobutene macromolecules have terminal double bonds to an extent of at least 50 mol%.

Claim 19 (Currently Amended): The mixture method according to claim 17, wherein

said polyisobutene block has been formed from polyisobutene macromolecules of which at

least 60 mol% based on the total number of polyisobutene macromolecules, have terminal

double bonds.

2

Claim 20 (Currently Amended): The mixture method according to claim 17, wherein said polyisobutene block has a number-average molecular weight Mn in the range from 200 to 20 000 daltons.

Claim 21 (Currently Amended): The mixture method according to claim 17, wherein said polyisobutene block has a polydispersity index (PDI) in the range from 1.05 to 10.

Claim 22 (Currently Amended): The mixture method according to claim 17, wherein one or more of said hydrophilic subunits (B) have been formed from repeat ethylene oxide units or ethylene oxide/propylene oxide units.

Claim 23 (Currently Amended): The mixture method according to claim 17, wherein one or more of said hydrophilic subunits (B) have been formed from monomer units selected from the group consisting of (meth)acrylic acid, including partly or fully neutralized (meth)acrylic acid, (meth)acrylates, vinyl acetate, vinyl alcohol, vinylpyrrolidone, allyl alcohol, styrene and hydrophilic derivatives of the listed monomer units, or from mixtures thereof.

Claim 24 (Currently Amended): The mixture method according to claim 17, wherein said polyisobutene block is functionalized by the introduction of polar groups and the functionalized polyisobutene block is optionally modified further.

Claim 25 (Currently Amended): The mixture method according to claim 24, wherein the functionalization of the polyisobutene block is carried out by a reaction which is selected from the group consisting of:

- i) a reaction with aromatic hydroxyl compounds in the presence of an alkylation catalyst to obtain aromatic hydroxyl compounds alkylated with polyisobutenes,
- ii) a reaction of the polyisobutene block with a peroxy compound to obtain an epoxidized polyisobutene,
- iii) a reaction of the polyisobutene block with an alkene which has an electron-poor double bond (enophile) in an ene reaction,
- iv) a reaction of the polyisobutene block with carbon monoxide and hydrogen in the presence of a hydroformylation catalyst to obtain a hydroformylated polyisobutene,
- v) a reaction of the polyisobutene block with a phosphorus halide or a phosphorus oxychloride to obtain a polyisobutene functionalized with phosphone groups,
- vi) a reaction of the polyisobutene block with a borane and subsequent oxidative cleavage to obtain a hydroxylated polyisobutene,
- vii) a reaction of the polyisobutene block with an SO3 source, preferably acetyl sulfate, to obtain a polyisobutene with terminal sulfonic acid groups, and
- viii) a reaction of the polyisobutene block with nitrogen oxides and subsequent hydrogenation to obtain a polyisobutene with terminal amino groups.

Claim 26 (Currently Amended): The $\frac{method}{method}$ according to claim 17, wherein said cosurfactant has an A_pB_q structure where p and q are each independently an integer from 1 to 8, or a comb structure composed of A and B.

Claim 27 (Currently Amended): The mixture method according to claim 17, wherein said surfactant is a surfactant with narrow homolog distribution.

Claim 28 (Currently Amended): The method of using a mixture according to claim 17, wherein the emulsion is a microemulsion for stabilizing emulsions and microemulsions.

Claim 29 (Currently Amended): The method according to claim 28, A microemulsion comprising a mixture of a surfactant and a cosurfactant according to claim 17, wherein said surfactant is a surfactant with narrow homolog distribution or one obtained under DMC catalysis.

Claim 30-33 (Canceled).

Claim 34 (Currently Amended): A detergent, cleaner, wetting agent, coating, adhesive, leather degreasing composition, humectant or textile treatment composition or a pharmaceutical, crop protection or cosmetic formulation, selected from the group consisting of sunscreen, skincare and hair styling composition, shower gel, shampoo, bath additive and scent oil, comprising, as well as customary ingredients, a mixture an emulsion stabilized according to claim 17.